

Superstars and Giant Gravitons in M-Theory

Frédéric Leblond¹, Robert C. Myers² and David C. Page³

^{1,2}Department of Physics, McGill University
3600 University Street, Montréal, Québec H3A 2T8, Canada

²Perimeter Institute for Theoretical Physics
35 King Street North, Waterloo, Ontario N2J 2W9, Canada

²Department of Physics, University of Waterloo
Waterloo, Ontario N2L 3G1, Canada

³Centre for Particle Theory
Department of Mathematical Sciences
University of Durham, Durham DH1 3LE, U.K.

ABSTRACT

Following hep-th/0109127, we show that a certain class of BPS naked singularities (superstars) found in compactifications of M-theory can be interpreted as being composed of giant gravitons. More specifically, we study superstars which are asymptotically $AdS_7 \times S^4$ and $AdS_4 \times S^7$ and show that these field configurations can be interpreted as being sourced by continuous distributions of spherical M2- and M5-branes, respectively, which carry internal momenta and have expanded on the spherical component of the space-time.

¹E-mail: fleblond@hep.physics.mcgill.ca

²E-mail: rcm@hep.physics.mcgill.ca

³E-mail: d.c.page@durham.ac.uk

1 Introduction

In a recent paper[1], convincing evidence was provided showing that certain supersymmetric naked singularities, appearing in type IIB supergravity compactified on $AdS_5 \times S^5$, can be interpreted in terms of distributions of giant gravitons[2]. These ‘superstar’ solutions correspond to the supersymmetric limit of a certain family of black holes. The latter were originally found when considering a consistent truncation of the type IIB supergravity theory dimensionally reduced to AdS_5 where they appear as charged black hole solutions. Once lifted back to the full ten-dimensional theory, the solutions carry internal momentum along the three commuting Killing angles on the five-sphere, and the supersymmetric limit still leaves a naked singularity. However, within the context of ten-dimensional type IIB superstring theory, there is a physical interpretation in which the singularities are generated by a distribution of giant gravitons, *i.e.*, an ensemble of spherical D3-branes which carry internal angular momentum and have expanded on the five-sphere[2]. This result was determined by examining the dipole field excited in the Ramond–Ramond five-form near the singularity of the supergravity solutions.

In this note, we study the analogous superstar solutions in M-theory compactified on $AdS_7 \times S^4$ and $AdS_4 \times S^7$. We show that these eleven-dimensional supergravity solutions can also be interpreted as being sourced by distributions of giant gravitons. Hence in this case, the constituent degrees of freedom are M2 and M5-branes, respectively. In the second part of this note, we study the behavior of test-brane probes in the background geometry of the M-theory superstars. By considering giant graviton probes, we are able to confirm the expansion of these configurations. We conclude with a brief discussion of our results. We consider dual giant graviton[3, 4] probes of superstars in an appendix.

2 Superstars in $AdS_7 \times S^4$

Eleven-dimensional supergravity appears as a particular low energy limit of M-theory[5]. The bosonic sector of this theory is composed of the graviton and the four-form field strength $F_{(4)}$. The latter allows for a spontaneous compactification[6] of the theory on $AdS_7 \times S^4$. Further in this background, there is a consistent truncation of the full theory to $\mathcal{N} = 4$ gauged $SO(5)$ supergravity in seven dimensions[7, 8]. In this note, we focus on solutions of a specific $\mathcal{N} = 2$ truncation of this theory where only the following seven-dimensional fields are retained: the metric, two scalars labelled by X_i ($i = 1, 2$) and two one-form gauge fields $A_{(1)}^i$ associated with the $U(1)^2$ Cartan subgroup of $SO(5)$ (see, *e.g.*, refs. [9, 10] for details). The correct Kaluza–Klein S^4 reduction ansatz, which is going to be used to lift seven-dimensional solutions to solutions of the full eleven-dimensional supergravity equations of motion, is[9, 10]

$$ds_{11}^2 = \tilde{\Delta}^{1/3} ds_7^2 + \frac{L^2}{\tilde{\Delta}^{2/3}} \left(\frac{1}{X_0} d\mu_0^2 + \sum_{i=1}^2 \frac{1}{X_i} \left[d\mu_i^2 + \mu_i^2 \left(d\phi_i + A_{(1)}^i/L \right)^2 \right] \right) \quad (1)$$

for the metric, and

$$\begin{aligned} *F_{(4)} = & -\frac{2}{L} \sum_{\alpha=0}^2 \left(X_\alpha^2 \mu_\alpha^2 - \tilde{\Delta} X_\alpha \right) \epsilon_{(7)} - \frac{1}{L} \tilde{\Delta} X_0 \epsilon_{(7)} - \frac{L}{2} \sum_{\alpha=0}^2 \frac{1}{X_\alpha} \overline{*} dX_\alpha \wedge d\mu_\alpha^2 \\ & - \frac{L^2}{2} \sum_{i=1}^2 \frac{1}{X_i^2} d\mu_i^2 \wedge \left(d\phi_i + A_{(1)}^i/L \right) \wedge \overline{*} F_{(2)}^i, \end{aligned} \quad (2)$$

for the field strength of the supergravity three-form. The internal four-sphere is parameterized by four angular variables: the ϕ_i 's are azimuthal (Killing) angles and the other angles are defined through the direction-cosines

$$\mu_0 = \sin \theta_1 \sin \theta_2 \quad \mu_1 = \cos \theta_1 \quad \mu_2 = \sin \theta_1 \cos \theta_2 , \quad (3)$$

such that $\mu_0^2 + \mu_1^2 + \mu_2^2 = 1$. The $\bar{*}$ denotes the Hodge dual with respect to the seven-dimensional metric ds_7^2 . Finally, L is the (asymptotic) radius of curvature of the four-sphere, $X_0 \equiv (X_1 X_2)^{-1}$ and $\tilde{\Delta} = \sum_{\alpha=0}^2 X_\alpha \mu_\alpha^2$.

The seven-dimensional $\mathcal{N} = 2$ supergravity admits doubly-charged AdS black hole solutions of the form[11],

$$ds_7^2 = -\frac{f}{(H_1 H_2)^{4/5}} dt^2 + (H_1 H_2)^{1/5} \left(\frac{1}{f} dr^2 + r^2 d\Omega_5^2 \right), \quad (4)$$

$$A_{(1)}^i = \left(\frac{1}{H_i} - 1 \right) dt \quad i = 1, 2 , \quad (5)$$

$$X_i = \frac{(H_1 H_2)^{2/5}}{H_i}, \quad (6)$$

where we have introduced

$$f = 1 - \frac{\mu}{r^4} + \frac{r^2}{4L^2} H_1 H_2, \quad (7)$$

$$H_i = 1 + \frac{q_i}{r^4}, \quad (8)$$

with μ a mass parameter and the q_i 's related to the physical charges of the black hole. We will use the following expression for the metric on the five-dimensional unit sphere,

$$d\Omega_5^2 = d\alpha_1^2 + \sin^2 \alpha_1 \left(d\alpha_2^2 + \sin^2 \alpha_2 \left[d\alpha_3^2 + \sin^2 \alpha_3 \left(d\alpha_4^2 + \sin^2 \alpha_4 d\alpha_5^2 \right) \right] \right). \quad (9)$$

The mass of this AdS_7 black hole is[12]

$$M = \frac{\pi^2}{4G_7} \left(\frac{5}{4} \mu + \sum_i q_i \right), \quad (10)$$

a quantity that will later be compared with the energy of the suggested ensemble of source branes.

The horizon structure of both the metric eq. (4) and its lift to eleven-dimensional supergravity using ansatz eq. (1) are essentially the same as can be seen by analysing the (rr) component of the corresponding metrics. The supersymmetric limit corresponds to $\mu = 0$ where the event horizon vanishes to reveal a naked singularity. Hence we denote these BPS configurations as ‘superstars.’ For $\mu = 0$, the entropy vanishes which means that the system should have a very simple physical interpretation in terms of a single arrangement of fundamental degrees of freedom. It is natural to conjecture that these degrees of freedom are giant gravitons, just as in the type IIB case[1]. The $\mu \neq 0$ black holes of finite entropy are likely still related to a similar ensemble of expanded branes but non-trivial interactions in the non-extremal case complicate the analysis. Hence we restrict ourselves to studying the BPS objects.

Following ref. [1] then, we make the conjecture that the superstars can be given a reasonable physical interpretation as the external fields around an ensemble of giant gravitons. The first evidence of this conjecture is simply that the off-diagonal form of the eleven-dimensional metric

(1) shows that the superstars carry internal momentum along the ϕ_i directions, just as giant gravitons would. However if the q_i 's are to be macroscopic distance scales (much larger than the eleven-dimensional Planck scale), then the mass of the superstar must be significantly larger than the energy of an individual giant graviton. This fact suggests that superstars must be composed of a collection of these fundamental objects.

For the detailed analysis, we begin for simplicity by considering superstars that are singly charged: $q_1 \neq 0$ and $q_2 = 0$. Since the source M2-branes must carry internal momentum along ϕ_1 , they should span the two-sphere parameterized by θ_2 and ϕ_2 . The collection of giant gravitons each with varying momenta should be distributed along the θ_1 -direction. Using a test-brane analysis[2], one finds that for a given internal momentum P_{ϕ_1} (the canonical conjugate to ϕ_1), a spherical M2-brane configuration on $AdS_7 \times S^4$ has two minima, one at $\sin \theta_1 = 0$ and the other at $\sin \theta_1 = P_{\phi_1}/N$, where N is the number of four-form flux quanta on the four-sphere. The former corresponds to a point-like graviton with angular momentum P_{ϕ_1} , and the latter to the giant graviton, with the same angular momentum, that has expanded to a fixed size on a two-dimensional submanifold of the S^4 . The point-like and giant gravitons preserve the same number of supersymmetries[3, 4] and have the same energy[2]:

$$E = \frac{P_{\phi_1}}{L} = \frac{N}{L} \sin \theta_1. \quad (11)$$

It is natural to assume that the superstar source must correspond to the expanded giant gravitons as the singularity in the metric extends over the entire four-sphere at $r = 0$.

A M2-brane is electrically charged with respect to the supergravity three-form potential. However, the spherical M2-branes will carry no net charge, but they will locally excite this field. If one considers a small (seven-dimensional) surface that encloses a portion of the sphere, one will find a net flux proportional to the number of M2-branes enclosed. In fact,

$$\int_{M_7} *F_4 = 16\pi G_{11} T_{M2} n_1, \quad (12)$$

where G_{11} is the eleven-dimensional gravitational constant, T_{M2} the tension of a M2-brane and n_1 the total number of giant gravitons enclosed. We are using the following conventions,

$$G_{11} = 16\pi^7 l_P^9, \quad G_7 = \frac{G_{11}}{V_{S^4}} = \frac{3}{8\pi^2 L^4} G_{11}, \quad T_{M2} = \frac{1}{4\pi^2 l_P^3}, \quad (13)$$

where l_P is the fundamental eleven-dimensional Planck length. Given the orientation of the giant gravitons described above, the surface M_7 is a manifold in the subspace spanned by $\{\theta_1, \phi_1, \alpha_i\}$. If we wish to consider the entire distribution, we naturally fix r and integrate over the remaining angles[1]. The ϕ_1 integration is of course trivial because the supergravity solution is smeared along this direction. In evaluating the expression eq. (12), we find that the only term in eq. (2) which gives a nonvanishing contribution is

$$-\frac{L^2}{2X_1^2} d\mu_1^2 \wedge d\phi_1 \wedge *F_{(2)}, \quad (14)$$

where

$$[*F_{(2)}]_{\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5} = -\frac{4q_1}{H_1^{6/5}} \sin^4 \alpha_1 \sin^3 \alpha_2 \sin^2 \alpha_3 \sin \alpha_4. \quad (15)$$

Hence the total number of giant gravitons n_1 is found to be

$$n_1 = \frac{q_1 L^2}{4\pi G_{11} T_{M2}} \int d\theta_1 d\phi_1 d^5\alpha_i \cos\theta_1 \sin\theta_1 \sin^4\alpha_1 \sin^3\alpha_2 \sin^2\alpha_3 \sin\alpha_4. \quad (16)$$

Dropping the θ_1 integration (along which the giant gravitons are assumed to be distributed) from eq. (16) and integrating over the other angles we find an expression for the density of giant gravitons as a function of θ_1 ,

$$\frac{dn_1}{d\theta_1} = \frac{q_1 N^2}{8L^4} \cos\theta_1 \sin\theta_1. \quad (17)$$

Given that we have detected a distribution of ‘electric dipole’ sources for $F_{(4)}$ at the singularity is certainly evidence of the presence of giant gravitons.

Further support for this result comes by considering the energy of the above distribution[1]. The test-brane analysis[2] found that the size and internal momentum of an individual giant graviton were related by: $P_{\phi_1} = N \sin\theta_1$. Combining this result with eq. (17), we find the total angular momentum of the distribution to be

$$\overline{P}_{\phi_1} = N \int_0^{\pi/2} d\theta_1 \sin\theta_1 \frac{dn_1}{d\theta_1} = \frac{1}{24} \frac{q_1 N^3}{L^4}. \quad (18)$$

According to eq. (11), the total energy is then

$$\overline{E} = \frac{1}{24} \frac{q_1 N^3}{L^5}. \quad (19)$$

The mass of the superstar (10), calculated by conventional supergravity means, is

$$M = \frac{\pi^2 q_1}{4G_7}. \quad (20)$$

Using eqs. (13) to write this mass in terms of N and L , one shows that it agrees exactly with the total energy eq. (19) of a distribution of giant gravitons. As it stands this result is somewhat of a curiosity. It is not clear why the result for the size of the giant graviton derived from a test-brane propagating in the $AdS_7 \times S^4$ background should apply here. However, we will find supporting evidence for this fact from our probe analysis in section 4.

It is straightforward to generalize the calculation to the case of superstars with two non-vanishing charges: $q_1 \neq 0$ and $q_2 \neq 0$. We consider the following embedding of S^4 in $\mathbf{R}^5$ with coordinates $x^{0,1,2,3,4}$,

$$x^0 = L\mu_0, \quad x^{2i-1} = L\mu_i \cos\phi_i, \quad x^{2i} = L\mu_i \sin\phi_i, \quad (21)$$

where $i = 1, 2$. Consequently, a giant graviton moving along ϕ_i has a radius

$$\rho_i = L\sqrt{1 - \mu_i^2}. \quad (22)$$

In analogy with the single charge calculation, the density of gravitons of a certain radius involves the $*F^{(4)}$,

$$\left[*F^{(4)} \right]_{\rho_i \phi_i \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5} = \frac{d\mu_i^2}{d\rho_i} \left[*F^{(4)} \right]_{\mu_i^2 \phi_i \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5} = 4q_i \rho_i \sin^4\alpha_1 \sin^3\alpha_2 \sin^2\alpha_3 \sin\alpha_4. \quad (23)$$

The corresponding density of giant gravitons for each direction is then

$$\frac{dn_i}{d\rho_i} = \frac{q_i N^2 \rho_i}{L^6}. \quad (24)$$

Treating this as two independent distributions, it is found that the total angular momentum carried by each set of giant gravitons is

$$\overline{P}_{\phi_i} = \frac{1}{24} \frac{q_i N^3}{L^4}. \quad (25)$$

These results match the total angular momentum calculated for the superstar solution, and we have complete agreement between the BPS mass of the superstar (10) and the total energy of the giant gravitons, $\overline{E} = \sum \overline{P}_{\phi_i}/L$.

3 Superstars in $AdS_4 \times S^7$

In this section we consider superstars which are asymptotically $AdS_4 \times S^7$. Their description is hypothesized to be in terms of M5-branes carrying internal momentum that have expanded on the S^7 . Eleven-dimensional supergravity admits a spontaneous compactification on $AdS_4 \times S^7$. The Kaluza–Klein reduction of this theory on S^7 leads to $\mathcal{N} = 8$ gauged $SO(8)$ supergravity in four dimensions. We focus on a specific $\mathcal{N} = 2$ truncation where only the following fields are retained: the metric, four scalars labelled by X_i ($i = 1, 2, 3, 4$) ($X_1 X_2 X_3 X_4 = 1$) and four one-form gauge fields $A_{(1)}^i$ (see, *e.g.*, refs. [9, 10] for details). The correct Kaluza–Klein S^7 reduction ansatz, which is going to be used to lift four-dimensional solutions to solutions of the full eleven-dimensional supergravity equations of motion, is [9, 10]

$$ds_{11}^2 = \tilde{\Delta}^{2/3} ds_4^2 + \frac{L^2}{\tilde{\Delta}^{1/3}} \sum_{i=1}^4 \frac{1}{X_i} \left(d\mu_i^2 + \mu_i^2 \left(d\phi_i + A_{(1)}^i/L \right)^2 \right) \quad (26)$$

for the metric, and

$$F_{(4)} = \frac{2}{L} \sum_{i=1}^4 \left(X_i^2 \mu_i^2 - \tilde{\Delta} X_i \right) \epsilon_{(4)} + \frac{L}{2} \sum_{i=1}^4 \frac{1}{X_i} \bar{*} dX_i \wedge d\mu_i^2 - \frac{L^2}{2} \sum_{i=1}^4 \frac{1}{X_i^2} d\mu_i^2 \wedge \left(d\phi_i + A_{(1)}^i/L \right) \wedge \bar{*} F_{(2)}^i, \quad (27)$$

for the field strength of the supergravity three-form. The space transverse to ds_4^2 is characterized by seven angular variables: the ϕ_i 's are azimuthal (Killing) angles and the other angular variables are defined through the direction-cosines

$$\mu_1 = \cos \theta_1 \quad \mu_2 = \sin \theta_1 \cos \theta_2 \quad \mu_3 = \sin \theta_1 \sin \theta_2 \sin \theta_3 \quad \mu_4 = \sin \theta_1 \cos \theta_2 \cos \theta_3, \quad (28)$$

such that $\mu_1^2 + \mu_2^2 + \mu_3^2 + \mu_4^2 = 1$. Finally, L is the (asymptotic) radius of curvature of the seven-sphere and $\tilde{\Delta} = \sum_{i=1}^4 X_i \mu_i^2$.

The corresponding four-dimensional action for the graviton, the scalars X^i and the one-form fields $A_{(1)}^i$ admit a quadruply charged AdS black hole solution [13, 14],

$$ds_4^2 = -\frac{f}{(H_1 H_2 H_3 H_4)^{1/2}} dt^2 + (H_1 H_2 H_3 H_4)^{1/2} \left(\frac{1}{f} dr^2 + r^2 d\Omega_2^2 \right), \quad (29)$$

$$A_{(1)}^i = \left(\frac{1}{H_i} - 1 \right) dt \quad i = 1, 2, 3, 4, \quad (30)$$

$$X_i = \frac{(H_1 H_2 H_3 H_4)^{1/4}}{H_i}, \quad (31)$$

where we have introduced

$$f = 1 - \frac{\mu}{r} + \frac{4r^2}{L^2} H_1 H_2 H_3 H_4, \quad (32)$$

$$H_i = 1 + \frac{q_i}{r}, \quad (33)$$

with μ a mass parameter and the q_i 's related to the physical charges of the black hole. We use the following expression for the metric on the two-dimensional unit sphere,

$$d\Omega_2^2 = d\alpha_1^2 + \sin^2 \alpha_1 d\alpha_2^2. \quad (34)$$

The mass of this AdS_4 black hole is [12]

$$M = \frac{1}{4G_4} \left(2\mu + \sum_i q_i \right), \quad (35)$$

a quantity that is to be compared with the energy of the hypothesized equivalent system composed of spherical M5-branes. Once again, the horizon structure of the lower dimensional realisation of the black hole and its lift to eleven-dimensional supergravity using ansatz eq. (26) are essentially the same. We assume that

$$q_1 \geq q_2 \geq q_3 \geq q_4. \quad (36)$$

For $q_1 \neq 0$ with all other charges vanishing, we find that for $\mu = 0$ there is no horizon behind which the curvature singularity at $r = 0$ is hidden. Whenever two charges or more have a finite value, there is a critical value for the non-extremality parameter, $\mu = \mu_{crit}$, corresponding to disappearing horizons. In other words, for $\mu > \mu_{crit}$ the geometry corresponds to a regular black hole and for $\mu < \mu_{crit}$, it corresponds to a naked singularity *i.e.*, to the so-called superstars. In any case, a generic feature is that for $\mu = 0$ (BPS limit) the area of the spherical horizon is zero.

Following the discussion of section 2, we consider superstars that are singly charged ($q_1 \neq 0$ and $q_2 = q_3 = q_4 = 0$) and assume that they are composed of M5-branes spanning a five-sphere parametrized by θ_2, θ_3 and ϕ_2, ϕ_3, ϕ_4 . The giant gravitons have non-zero internal momentum along ϕ_1 , are distributed along the θ_1 -direction and have an energy given by [2]

$$E = \frac{P_{\phi_1}}{L} = \frac{N}{L} \sin^4 \theta_1, \quad (37)$$

where N is the number of seven-form (the dual of $F^{(4)}$) flux quanta on the seven-sphere. A M5-brane is magnetically charged with respect to the supergravity three-form potential. The spherical M5-branes carry no net charge but nevertheless excite the field locally. If one considers a small surface enclosing a portion of the sphere, one finds a net flux proportional to the number of M5-branes enclosed,

$$\int_{M_4} F_4 = 16\pi G_{11} T_{M5} n_1, \quad (38)$$

where G_{11} is the eleven-dimensional gravitational constant, T_{M5} is the tension of a M5-brane and n_1 is the total number of giant gravitons making the superstar. Given the orientation of the giant gravitons described above, M_4 is a manifold in the subspace spanned by $\{\theta_1, \phi_1, \alpha_i\}$. The ϕ_1 integration is trivial because the supergravity solution is smeared along that direction. We are using the following conventions,

$$G_{11} = 16\pi^7 l_P^9, \quad G_4 = \frac{G_{11}}{V_{S^7}} = \frac{3}{\pi^4 L^7} G_{11}, \quad T_{M5} = \frac{1}{2^5 \pi^5 l_P^6}, \quad (39)$$

where l_P is the fundamental eleven-dimensional Planck length. In evaluating the expression eq. (38) we find that all except the last term of eq. (27) lead to a vanishing integral. The following term of $F_{(4)}$ is therefore the only one that is relevant to the present analysis,

$$F_{(4)} = -\frac{L^2}{2} \frac{1}{X} d(\mu_1^2) \wedge d\phi_1 \wedge \bar{*}F_{(2)}, \quad (40)$$

where

$$\left[\bar{*}F_{(2)}\right]_{\alpha_1\alpha_2} = -\frac{q_1}{H^{3/2}} \sin \alpha_1. \quad (41)$$

In the end, the total number of giant gravitons n_1 is found to be

$$n_1 = \frac{q_1 L^2}{4\pi G_{11} T_{M5}} \int d\theta_1 d\phi_1 d\alpha_1 d\alpha_2 \cos \theta_1 \sin \theta_1 \sin \alpha_1. \quad (42)$$

Dropping the θ_1 integration (along which the giant gravitons are assumed to be distributed) from eq. (42) and integrating over the other angles we find an expression for the density of giant gravitons as a function θ_1 ,

$$\frac{dn_1}{d\theta_1} = \frac{8q_1 N^{1/2}}{\sqrt{2}L} \cos \theta_1 \sin \theta_1. \quad (43)$$

As in section 2, we have detected a distribution of 'electric dipole' sources for $F^{(4)}$ at the singularity, which is certainly evidence of the presence of giant gravitons.

In order to give further support for this result, we calculate the energy of the above distribution. A test-brane analysis showed[2] that a single giant graviton is associated with an internal momentum given by

$$P_{\phi_1} = N \sin^4 \theta_1. \quad (44)$$

Combining this result with eq. (43), we find the total angular momentum of the distribution,

$$\bar{P}_{\phi_1} = N \int_0^{\pi/2} d\theta_1 \sin \theta_1 \frac{dn_1}{d\theta_1} = \frac{4\sqrt{2}}{3} \frac{q_1 N^{3/2}}{L}, \quad (45)$$

which corresponds, according to eq. (11), to a total energy

$$\bar{E} = \frac{4\sqrt{2}}{3} \frac{q_1 N^{3/2}}{L^2}. \quad (46)$$

The mass of the superstar (35), calculated by conventional supergravity means, is

$$M = \frac{q_1}{4G_4}. \quad (47)$$

Using eqs. (39) to write the mass of the superstar in terms of N and L , one shows that it agrees exactly with the total energy eq. (46) of a distribution of giant gravitons. This is strong evidence that the asymptotically $AdS_4 \times S^7$ superstars are in fact M-theory objects composed of microscopic spherical M5-branes that have expanded on the seven-sphere.

Following the procedure of the previous section, we can easily generalize to the case of superstars with none of the four charges vanishing: $q_i \neq 0$ ($i = 1, 2, 3, 4$). We consider the following embedding of S^7 in $\mathbf{R}^8$ with coordinates $x^{1,\dots,8}$,

$$x^{2i-1} = L\mu_i \cos \phi_i, \quad x^{2i} = L\mu_i \sin \phi_i. \quad (48)$$

Consequently, a giant graviton moving along ϕ_i has a radius of

$$\rho_i = L\sqrt{1 - \mu_i^2}. \quad (49)$$

In analogy with the single charge calculation, the density of gravitons of a certain radius involves the $F^{(4)}$,

$$F_{\rho_i \phi_i \alpha_1 \alpha_2}^{(4)} = \frac{d\mu_i^2}{d\rho_i} F_{\mu_i^2 \phi_i \alpha_1 \alpha_2}^{(4)} = q_i \rho_i \sin \alpha_1. \quad (50)$$

The corresponding density of giant gravitons for each direction is then

$$\frac{dn_i}{d\rho_i} = \frac{8q_i \sqrt{N} \rho_i}{\sqrt{2} L^3}. \quad (51)$$

Just as above, it is found that the total angular momentum carried by each set of giant gravitons is

$$\overline{P}_{\phi_i} = \frac{4\sqrt{2}}{3} \frac{q_i N^{3/2}}{L}. \quad (52)$$

These results correspond to the total angular momentum calculated for the superstar solution, and we have complete agreement between the BPS mass of the superstar (35) and the total energy of the giant gravitons, $\overline{E} = \sum \overline{P}_{\phi_i} / L$.

4 Giant graviton probes

In the previous sections, we showed that the two eleven-dimensional superstar geometries have an interpretation as the external fields for a collection of giant gravitons. That is, we should regard these configurations as solutions of the eleven-dimensional supergravity coupled to the brane actions of the corresponding giant gravitons. The primary evidence is the fact that the naked singularities behave as sources for the appropriate dipoles of the four-form field strength. However, to complete the analysis, we should also show that we also have a solution of the brane equations of motion. As a step in this direction, we consider a giant graviton probe in the superstar background. We will indeed find expanded configurations in the limit as $r \rightarrow 0$. Further for these configurations, we will confirm the result: $P_{\phi_1} = N \sin^{p-1} \theta_1$ (where $p+1$ is the dimension of the M-brane's worldvolume). It was with this result that we showed that the distribution of giant gravitons derived from flux considerations matched the total internal momentum of the superstar.

4.1 Probing superstars in $AdS_7 \times S^4$

In the following, we present a brief summary of the M2-brane probe calculations for the superstars presented in section 2. We will restrict our attention to the simplest case of a singly charged superstar (*i.e.* only q_1 is nonvanishing). The results of the doubly charged case are less clear — we will return to this point in the discussion.

Our giant graviton probe will be a spherical M2-brane inside the S^4 at constant θ_1 and moving on a circle in the ϕ_1 direction. We will first consider this configuration at a finite radius in the anti-de Sitter space (*i.e.* away from the singularity) and then consider taking the test-brane to $r = 0$. The M2-brane couples to the three form potential $A^{(3)}$ satisfying $dA^{(3)} = F^{(4)}$. Hence to evaluate the worldvolume action, we need to dualize $*F^{(4)}$ given in eq. (2) and integrate to find $A^{(3)}$. The result is

$$F^{(4)} = -\frac{1}{L}H_1\frac{(\Delta+2)}{\Delta^2}\sin^2\theta_1\cos\theta_1Ld\theta_1\wedge\sin\theta_2\cos\theta_2Ld\theta_2\bigwedge_j[Ld\phi_j+A_{(1)}^j] + \dots \quad (53)$$

$$= d\left(-H_1\frac{\sin^3\theta_1}{\Delta}\wedge\sin\theta_2\cos\theta_2Ld\theta_2\bigwedge_j[Ld\phi_j+A_{(1)}^j]\right) = dA^{(3)}. \quad (54)$$

The probe action is then

$$S_2 = -T_{M2}\int dt d\theta_2 d\phi_2 \left[\sqrt{-g} - A_{t\theta_2\phi_2}^{(6)} - \dot{\phi}_1 A_{\phi_1\theta_2\phi_2}^{(4)}\right] = \frac{N}{L}\int dt \left[-\Delta^{-1/2}\sin^2\theta_1\left(\frac{f}{H_1} - \Delta^{-1}H_1\mu_1^2[L\dot{\phi}_1 + (H_1^{-1}-1)]^2\right)^{1/2} + \frac{H_1}{\Delta}\sin^3\theta_1(L\dot{\phi}_1 + H_1^{-1}-1)\right], \quad (55)$$

Now fixing the momentum $p \equiv P_{\phi_1}/N$, we find the Hamiltonian

$$\mathcal{H} = \frac{N}{L}\left[\sqrt{\frac{f}{H_1}\left(\frac{p^2}{H_1} + \tan^2\theta_1(p - \sin\theta_1)^2\right)} + (1 - H_1^{-1})p\right]. \quad (56)$$

Minimizing $\mathcal{H}(r, \theta_1)$ with respect to θ_1 yields $\theta_1 = 0$ or $\sin\theta_1 = p$ for arbitrary values of r . However, to find a true solution of the equations of motion, we must also minimize with respect to the radius. Setting $\sin\theta_1 = p$, a short calculation shows that the minimum energy configuration is at $r=0$, and that the energy of this configuration satisfies the BPS relation:

$$\mathcal{H}(r=0, \sin\theta_1=p) = \frac{N}{L}p = \frac{P_{\phi_1}}{L}. \quad (57)$$

4.2 Probing superstars in $AdS_4 \times S^7$

The four-form and metric corresponding to the superstars in $AdS_4 \times S^7$ were given in section 3. The probe which we shall be studying in this case is a spherical M5-brane inside the S^7 at constant θ_1 and moving on a circle in the ϕ_1 direction. Again, we begin by placing the

probe away from the singularity and consider the limit as $r \rightarrow 0$. The M5-brane couples to the six-form potential $A^{(6)}$ satisfying $dA^{(6)} = *F^{(4)}$ and so our first task is to dualize $F^{(4)}$ given in eq. (27) and integrate to find $A^{(6)}$. We find

$$\begin{aligned} *F^{(4)} &= -\frac{2}{L}H_1\frac{(2\Delta+1)}{\Delta^2}\sin^5\theta_1\cos\theta_1Ld\theta_1\wedge\sin^3\theta_2\cos\theta_2Ld\theta_2 \\ &\quad \wedge\sin\theta_3\cos\theta_3Ld\theta_3\bigwedge_j[Ld\phi_j+A_{(1)}^j]+\dots \end{aligned} \quad (58)$$

$$\begin{aligned} &= d\left(-H_1\frac{\sin^6\theta_1}{\Delta}\wedge\sin^3\theta_2\cos\theta_2Ld\theta_2\wedge\sin\theta_3\cos\theta_3Ld\theta_3\bigwedge_j[Ld\phi_j+A_{(1)}^j]\right) \\ &= dA^{(6)}. \end{aligned} \quad (59)$$

Evaluating the probe action for the configuration described above yields

$$\begin{aligned} S_5 &= -T_{M5}\int dt d\theta_2 d\theta_3 d\phi_2 d\phi_3 d\phi_4 \left[\sqrt{-g} + A_{t\theta_2\theta_3\phi_2\phi_3\phi_4}^{(6)} + \dot{\phi}_1 A_{\phi_1\theta_2\theta_3\phi_2\phi_3\phi_4}^{(4)}\right] \\ &= \frac{N}{L}\int dt \left[-\Delta^{-1/2}\sin^5\theta_1\left(\frac{f}{H_1} - \Delta^{-1}H_1\mu_1^2[L\dot{\phi}_1 + (H_1^{-1}-1)]^2\right)^{1/2}\right. \\ &\quad \left. + \frac{H_1}{\Delta}\sin^6\theta_1(L\dot{\phi}_1 + H_1^{-1}-1)\right]. \end{aligned} \quad (60)$$

Again fixing $p \equiv P_{\phi_1}/N$, we find the Hamiltonian

$$\mathcal{H} = \frac{N}{L}\left[\sqrt{\frac{f}{H_1}\left(\frac{p^2}{H_1} + \tan^2\theta_1(p - \sin^4\theta_1)^2\right)} + (1 - H_1^{-1})p\right]. \quad (61)$$

Extremizing $\mathcal{H}$ with respect to θ_1 yields minima at $\theta_1 = 0$ and $\sin^4\theta_1 = p$ independent of the radius. The r -dependence of the Hamiltonian is essentially as before and we once again find a solution to the equations of motion at $r = 0$ satisfying the BPS relation :

$$\mathcal{H}(r=0, \sin^4\theta_1=p) = \frac{P_{\phi_1}}{L}. \quad (62)$$

5 Discussion

In this note, we have extended the analysis of ref. [1] to the context of M-theory. In particular, we have studied superstar solutions of eleven-dimensional supergravity in asymptotically $AdS_7 \times S^4$ and $AdS_4 \times S^7$ space-times. While these BPS configurations contain naked singularities, we have argued that M-theory provides a natural external source in the form of an ensemble of giant gravitons. Hence we should regard the superstars as solutions of the combined equations of motion of eleven-dimensional supergravity coupled to the worldvolume actions of the corresponding giant gravitons.

In section 4, we considered the dynamics of giant gravitons in the superstar backgrounds. With this analysis, we were able to recover the result that the expanded branes satisfy:

$$P_{\phi_1} = N \sin^{p-1}\theta_1 \quad (63)$$

where $p+1$ is the dimension of the M-brane's worldvolume. We have also confirmed that this result applies for a giant graviton probe in the type IIB superstar background considered

in ref. [1]. It is curious that these relations between the internal momentum of the various giant gravitons and their expansion in the internal space is identical to that derived in the corresponding $AdS_m \times S^n$ background. Certainly the energy–momentum relations for the BPS configurations must hold in the superstar backgrounds, as this is dictated by supersymmetry. However, the latter does not require the dynamical details such as the size to remain unperturbed in the superstar backgrounds. It would be interesting to investigate whether this result is simply a coincidence or whether there is a deeper reason why the expansion of the giant gravitons matches in these different backgrounds.

One should also question the validity of the result presented in eq. (63) since it relies on pushing past the expected regime of validity of the test–brane analysis. We expect, *e.g.*, that the usual Born–Infeld form of the action is only a leading order result that should be valid to describe the low energy dynamics of branes in slowly varying background fields. We must expect that there are higher derivative corrections that must become important in rapidly varying backgrounds, *e.g.*, in regions of strong curvature. For instance the leading order corrections to the worldvolume action can be systematically calculated in superstring theory [15]. In the present analysis, we begin by placing the giant gravitons away from the origin but then find that minimizing the energy requires that we set $r = 0$. However, this is the position of the naked singularity in the singly charged superstar, and so in fact, it is a location where the curvatures are diverging. The fact that we get a simple and consistent result seems to indicate that the higher curvature corrections must cancel out in the final analysis. However, clearly this point deserves a better understanding. In particular, one should note that the analysis in section 4 only considers the case of a singly charged superstar. We were unable to confirm the same results for the general case, but it may well be that strong curvature effects play an important role in these cases. Certainly, the analysis for the type IIB superstars [1] show the multiply charged systems to be more exotic.

Acknowledgements

Research by FL and RCM is supported in part by NSERC of Canada and Fonds FCAR du Québec. DCP is funded by the Engineering and Physical Sciences Research Council of Britain. FL would like to thank the Perimeter Institute for Theoretical Physics and the University of Waterloo’s Department of Physics for their kind hospitality during this work. We would like to thank Clifford Johnson for useful conversations and correspondence. DCP would also like to thank Ken Lovis and especially Douglas Smith for extensive conversations. While this paper was being completed, ref. [16] appeared which overlaps with the material presented in sections 2 and 3.

A Dual giant graviton probes

We have also considered probing the M–theory superstars with dual giant graviton probes, in analogy with the analysis presented in ref. [1].

A.1 Dual giants in $AdS_7 \times S^4$

In $AdS_7 \times S^4$, the dual giants are spherical M5-branes, spanning the S^5 of the AdS_7 space at constant r . They orbit on the S^4 at constant θ_1 and θ_2 with fixed angular momentum P_{ϕ_i}

conjugate to the angles ϕ_i .

The probe Lagrangian takes the following form (after integrating over the 5-sphere):

$$\begin{aligned} \mathcal{L} = & \frac{\tilde{N}}{L} \left[-\Delta^{1/2} \left(\frac{r}{\tilde{L}} \right)^5 \left(f \left(\mu_0^2 + \sum_i \frac{\mu_i^2}{H_i} \right) - \sum_i H_i \mu_i^2 [L\dot{\phi}_i + (H_i^{-1} - 1)]^2 \right)^{1/2} \right. \\ & \left. + \frac{r^6}{\tilde{L}^6} \Delta + \sum_i \frac{\mu_i^2 q_i}{\tilde{L}^4} (L\dot{\phi}_i - 1) \right] \end{aligned} \quad (64)$$

Here $\tilde{L} = 2L$ is the radius of the AdS space and $\tilde{N} = L\tilde{L}^2 A_2 T_2$, where A_2 is the area of a unit S^2 and T_2 is the tension of an M2 brane. Fixing the value of $\tilde{p}_i \equiv P_{\phi_i}/\tilde{N}$ we find the Hamiltonian:

$$\begin{aligned} \mathcal{H} = & \frac{\tilde{N}}{L} \left[\sqrt{f \left(\mu_0^2 + \sum_i \frac{\mu_i^2}{H_i} \right) \left(\sum_i \frac{(\tilde{p}_i - \frac{\mu_i^2 q_i}{\tilde{L}^4})^2}{H_i \mu_i^2} + \Delta \frac{r^{10}}{\tilde{L}^{10}} \right)} \right. \\ & \left. + \sum_i (1 - H_i^{-1}) \left(\tilde{p}_i - \frac{\mu_i^2 q_i}{\tilde{L}^4} \right) - \Delta \frac{r^6}{\tilde{L}^6} + \sum_i \frac{\mu_i^2 q_i}{\tilde{L}^4} \right] \end{aligned} \quad (65)$$

Mininima of this action saturating the BPS bound $\mathcal{H} = (N/L) \sum p_i$ occur whenever the following equations are satisfied:

$$\frac{r^4}{\tilde{L}^4} \mu_i^2 = \tilde{p}_i - \frac{q_i \mu_i^2}{\tilde{L}^4}. \quad (66)$$

Using the constraint that $\sum \mu_i^2 = 1$, one can reduce these equations to the same form as presented in eqs. (28) and (29) of ref. [1].

A.2 Dual giants in $AdS_4 \times S^7$

In $AdS_4 \times S^7$, the giant gravitons are M5-branes whereas the dual giants are spherical M2-branes, spanning the S^2 of the AdS_4 space at constant r . They orbit on the S^7 at constant θ_1 , θ_2 and θ_3 with fixed angular momentum P_{ϕ_i} conjugate to the angles ϕ_i .

The probe Lagrangian takes the following form (after integrating over the 2-sphere):

$$\begin{aligned} \mathcal{L} = & \frac{\tilde{N}}{L} \left[-\Delta^{1/2} \frac{r^2}{\tilde{L}^2} \left(\sum_i \frac{\mu_i^2}{H_i} f - \sum_i H_i \mu_i^2 [L\dot{\phi}_i + (H_i^{-1} - 1)]^2 \right)^{1/2} \right. \\ & \left. + \frac{r^3}{\tilde{L}^3} \Delta + \sum_i \frac{\mu_i^2 q_i}{\tilde{L}} (L\dot{\phi}_i - 1) \right] \end{aligned} \quad (67)$$

Here we have introduced $\tilde{L} = L/2$ as the radius of the AdS space and $\tilde{N} = L\tilde{L}^5 A_5 T_5$ where A_5 is the area of a unit S^5 and T_5 is the tension of an M5-brane. Fixing the value of $\tilde{p}_i \equiv P_{\phi_i}/\tilde{N}$ we find the Hamiltonian:

$$\begin{aligned} \mathcal{H} = & \frac{\tilde{N}}{L} \left[\sqrt{\left(\sum_i \frac{\mu_i^2}{H_i} f \right) \left(\sum_i \frac{(\tilde{p}_i - \frac{\mu_i^2 q_i}{\tilde{L}})^2}{H_i \mu_i^2} + \Delta \frac{r^4}{\tilde{L}^4} \right)} \right. \\ & \left. + \sum_i (1 - H_i^{-1}) \left(\tilde{p}_i - \frac{\mu_i^2 q_i}{\tilde{L}} \right) - \Delta \frac{r^3}{\tilde{L}^3} + \sum_i \frac{\mu_i^2 q_i}{\tilde{L}} \right] \end{aligned} \quad (68)$$

Mininima of this action saturating the BPS bound $\mathcal{H} = (N/L) \sum p_i$ occur whenever the following equations are satisfied:

$$\frac{r}{\tilde{L}} \mu_i^2 = \tilde{p}_i - \frac{q_i \mu_i^2}{\tilde{L}}. \quad (69)$$

References

- [1] R.C. Myers and O. Tafjord, *Superstars and Giant Gravitons*, hep-th/0109127, to appear in JHEP.
- [2] J. McGreevy, L. Susskind and N. Toumbas, JHEP **0006**, 008 (2000) [hep-th/0003075].
- [3] M.T. Grisaru, R.C. Myers and O. Tafjord, JHEP **0008**, 040 (2000) [hep-th/0008015].
- [4] A. Hashimoto, S. Hirano and N. Itzhaki, JHEP **0008**, 051 (2000) [hep-th/0008016].
- [5] E. Witten, Nucl. Phys. B **443**, 85 (1995) [hep-th/9503124].
- [6] P.G. Freund and M.A. Rubin, Phys. Lett. B **97**, 233 (1980).
- [7] H. Nastase, D. Vaman and P. van Nieuwenhuizen, Phys. Lett. B **469**, 96 (1999) [hep-th/9905075].
- [8] H. Nastase, D. Vaman and P. van Nieuwenhuizen, Nucl. Phys. B **581**, 179 (2000) [hep-th/9911238].
- [9] M. Cvetič, M.J. Duff, P. Hoxha, J.T. Liu, H. Lu, J.X. Lu, R. Martinez-Acosta, C.N. Pope, H. Sati, T.A. Tran, Nucl. Phys. B **558**, 96 (1999) [hep-th/9903214].
- [10] M.J. Duff, *TASI lectures on branes, black holes and anti-de Sitter space*, hep-th/9912164.
- [11] M. Cvetič and S.S. Gubser, JHEP **9904**, 024 (1999) [hep-th/9902195].
- [12] R.G. Cai, Phys. Rev. D **63**, 124018 (2001) [hep-th/0102113].
- [13] M.J. Duff and J.T. Liu, Nucl. Phys. B **554**, 237 (1999) [hep-th/9901149].
- [14] W.A. Sabra, Phys. Lett. B **458**, 36 (1999) [hep-th/9903143].
- [15] C.P. Bachas, P. Bain and M.B. Green, JHEP **9905**, 011 (1999) [hep-th/9903210].
- [16] V. Balasubramanian and A. Naqvi, *Giant Gravitons and a Correspondence Principle*, hep-th/0111163.